

# Homework 1 for ECON 21130

T. Lamadon

January 15, 2018

## 1 Probability and statistics

1. (10 points) Let  $X_1, \dots, X_n \sim \text{Uniform}(0,1)$  and let  $Y_n = \max X_1, \dots, X_n$ , what is  $E(Y_n)$  ?
2. (10 points) 2 players go in 2 different rooms where they each toss a coin simultaneously. Each player then has to guess the other player's toss outcome. The players can discuss their strategy beforehand, but cannot communicate once they enter their designated rooms. If both guessed wrong, they both get 0, if one or both correctly guessed the other players' toss outcome then both win a payoff of 1. Find the optimal strategy and the associated expected payoff per player (hint, it is strictly higher than 0.75).

## 2 OLS, endogeneity and conditioning

In this problem we are going to look at the effect of changing the conditioning set in a regression framework. Consider the following model for some data  $(Y_i, X_i, T_i)$ :

$$Y_i = \gamma T_i + \beta X_i + \epsilon_i$$

where  $T_i$  is a treatment of interest,  $X_i$  is a covariate,  $\epsilon_i$  is a residual and  $\beta \neq 0$ . Assume further that  $T_i, X_i, \epsilon_i$  are jointly normally distributed according to:

$$\begin{bmatrix} \epsilon_i \\ X_i \\ T_i \end{bmatrix} \sim N\left(0, \Omega\right) \quad \text{with} \quad \Omega = \begin{bmatrix} 1 & \rho_1 & 0 \\ \rho_1 & 1 & \rho_2 \\ 0 & \rho_2 & 1 \end{bmatrix}$$

We are interested in the parameter  $\gamma$  that measures the treatment effect.

1. (15 points) What is the joint distribution of  $(T_i, \epsilon_i)$  unconditional of  $X_i$ ? Does this tell us that  $T_i \perp \epsilon_i$ ? Does this mean that  $(T_i \perp \epsilon_i) | X_i$ ? Explain.

First we are interested in the result of the regression of  $Y_i$  on  $T_i$  only, excluding  $X_i$ . This requires us to express  $\mathbb{E}(Y_i | T_i)$ . We are going to use the result that if some variables  $X_{i,1}, X_{i,2}, X_{i,3}$  are joint normal then one can write each as linear combination of the two others. Hence there exist  $\alpha_1, \alpha_2$  such that  $X_{i,3} = \alpha_1 X_{i,1} + \alpha_2 X_{i,2} + u_i$  where  $u_i$  is a random variable, normally distributed and independent of  $X_{i,1}, X_{i,2}$ .

2. (10 points) Given that  $T_i$  and  $X_i$  are jointly normal, we can write  $X_i = aT_i + u_i$  for some scalar  $a$  and a normal random variable  $u_i$  independent of  $T_i$ . Use the entries of the  $\Omega$  matrix and express  $Cov(T_i, X_i)$  as a function of  $a$  to show that  $a = \rho_2$ .
3. (10 points) Use the previous expression for  $X_i$  to show that  $\mathbb{E}[Y_i|T_i] = \clubsuit \cdot T_i$ . Report the expression for  $\clubsuit$  as a function of the parameters  $\gamma, \beta, \rho_2$ .
4. (10 points) Under what condition does the regression coefficient  $\clubsuit$  provide an unbiased estimate of  $\gamma$ ? How can you interpret this condition as a condition on the relationship between  $X_i$  and  $T_i$ ?

We are now interested in the result of the regression of  $Y_i$  on  $T_i, X_i$  jointly. This requires us to express  $\mathbb{E}(Y_i|T_i, X_i)$ .

5. (15 points) Given that  $T_i, X_i$  and  $\epsilon_i$  are jointly normal, we can write  $\epsilon_i = b \cdot T_i + c \cdot X_i + v_i$  for some scalars  $b, c$  and a normal random variable  $v_i$  independent of  $T_i, X_i$ . Similar to the previous part, show that using  $Cov(X_i, \epsilon_i)$  and  $Cov(T_i, \epsilon_i)$  together with the entry of  $\Omega$  we can establish that:

$$\begin{aligned} 0 &= b + c\rho_2 \\ \rho_1 &= b\rho_2 + c \end{aligned}$$

leading to the following expression  $\mathbb{E}[\epsilon_i|T_i=t, X_i=x] = \frac{\rho_1}{1-\rho_2^2} \cdot x - \frac{\rho_1\rho_2}{1-\rho_2^2} \cdot t$  (which you are not asked to derive).

6. (10 points) We are then ready to find our regression expression of  $Y_i$  on  $T_i, X_i$ . To do so find the missing parts in the following equation:

$$\mathbb{E}[Y_i|X_i = x, T_i = t] = \blacksquare \cdot t + \blacklozenge \cdot x.$$

7. (10 points) Under what sufficient conditions does  $\blacksquare$  provide an unbiased estimate of  $\gamma$ ? How can you interpret this condition as a condition on the relationship between  $X_i$  and  $T_i$  or between  $\epsilon_i$  and  $X_i$ .
8. (10 points) [Bonus] If a researcher then tells you “It is always better to condition on all possible observables available to you”, in the light of the previous question, what would you respond? Is it possible for  $\blacksquare$  to be more strongly biased than  $\clubsuit$ ?